

DOI: 10.5281/zenodo.11042551

PHYSICAL EXERCISE IN NECK PAIN AND EMOTIONAL WELL-BEING IN CAREGIVER BURNOUT

Ana Díaz Cevallos^{1*}, Gyna Michelle Galarza Sánchez², José Luis Erazo Parra³, María Eugenia Lucena de Ustáriz⁴, Lisbeth Reales Chacón⁵

¹Universidad Nacional de Chimborazo, Ecuador, criss8diazcevallos@gmail.com, <https://orcid.org/0000-0003-0177-2734>

²Universidad Metropolitana, Quito, Ecuador. ggalarza@umet.edu.ec, <https://orcid.org/0009-0004-6755-7994>

³Universidad Nacional de Chimborazo, Ecuador. jlerazo@unach.edu.ec, <https://orcid.org/0000-0003-3149-0067>

⁴Universidad Nacional de Chimborazo, Ecuador. mlucena@unach.edu.ec, <https://orcid.org/0000-0001-9120-345X>

⁵Universidad Nacional de Chimborazo, Ecuador. lisbeth.reales@unach.edu.ec, <https://orcid.org/0000-0002-4242-3429>

Received: 11/11/2025

Accepted: 18/11/2025

Corresponding Author: Ana Díaz Cevallos
(criss8diazcevallos@gmail.com)

ABSTRACT

Most of the family members of people with Cerebral Palsy (CP) are those who function as informal caregivers. On certain occasions, their work affects their quality of life and their health, and can trigger lumbar and cervical pain. Studies carried out in several countries show that they are a highly vulnerable population, with poor coping strategies that favor their adaptive process. The present study aims to determine the effects of physical exercise on neck pain present in caregivers of people with CP. It is quasi-experimental quantitative research, with a pretest-post-test design with a control group, with a non-probabilistic sample due to accessibility considering the characteristics of the participants for the study. The research carried out the analysis of the functional effects on the cervical and emotional spine of the implementation of a physical exercise program in caregivers of people with CP. The intervention with a physical exercise program for 16 weeks in an experimental group of 100 informal CP caregivers for 19.83 ± 1.2 years improved caregiver burden, emotional health and neck pain. In contrast, the control group after 16 weeks worsened anxiety. The results of the present study collaborate as a non-pharmaceutical treatment for anxiety and neck pain, being the practice of physical exercise for 16 weeks a good option to alleviate the symptoms of Caregiver Overload Syndrome.

KEYWORDS: Caregivers, Cerebral Palsy, Physical Exercise.

1. INTRODUCTION

Cerebral Palsy (CP) is the physical disability that produces alterations in mobility and development, as well as others in sensation, perception, cognition, communication and behavior. (1). The family of subjects with CP tries to improve the quality of life of the people in their care, since a correlation is shown between the degree of motor impairment and the decrease in quality of life in the pediatric and adolescent population with CP. In addition to the quality of life of people with CP, the quality of life of their caregivers is also affected. Caregivers of a family member with CP often have health problems associated with the exercise of this work. Among the main diseases are: stress, headache, sleep disturbances, anxiety and neck pain. (2)(3). There is evidence that the use of supervised aerobic exercise, performed three times a week at moderate intensity for a minimum of eight weeks, has become an alternative treatment to medication in the management of depression and spinal pain. (4-6). Exercise increases blood flow, vascular density, during exercise the nervous system increases the levels of neurotransmitters such as: endorphins, norepinephrine, serotonin reducing anxiety, pain and improving mood and the feeling of well-being. (7,8). These benefits can also be seen reflected in caregivers of people with chronic neurological diseases, in different studies on the practice of physical exercise have shown positive effects mainly in reducing the caregiver's burden. (9,10). The physical and emotional health of the caregiver is important, and there is currently little scientific evidence on treatments, prevention and promotion measures that help improve the quality of life of this group of people, this was the main motivation for carrying out this research, which is why the objective of this study.

2. METHODS

2.1. Design

The present study is a quasi-experimental quantitative research, with a pre-test-post-test design with a control group, with a non-probabilistic sampling for accessibility considering the characteristics of the participants for the study. The research analyzed the physiological and emotional effects of the implementation of a physical exercise program in caregivers of people with CP.

2.2. Participants

A total of 200 women with a mean age of 48.25 ± 1.29 years, caregivers of a family member with CP,

who presented Caregiver Overload Syndrome, participated. The sample was divided into two groups: a so-called intervention group, which carried out a 16-week exercise program with a frequency of 3 times a week, and a control group whose participants followed their usual lives.

The inclusion criteria were: being a caregiver of a family member with CP, for a period between 18 and 25 years, with Caregiver Overload Syndrome, assessed with a score higher than 46 in the Zarit test, who did not have health problems that prevented them from practicing physical activity and who agreed to participate freely and voluntarily in the research. The exclusion criteria were being a caregiver for a family member with another dependent disease, presenting cardiovascular risk and/or pre-existing heart disease.

2.3. Procedure

The following variables were investigated: family functionality with the APGAR test, caregiver burden with the Zarit test, emotional well-being with the GHQ 28 questionnaire and neck pain with the Oswestry test. The physical exercise program was applied and supervised by a physiotherapy professional for a period of 16 weeks, 3 days a week (with a day off), for 60 minutes, the exercises were performed in a group manner. In the first weeks of the physical exercise program, it was started at a low intensity and a progressive increase was made until reaching a moderate intensity by week 16.

2.4. Statistical Analysis

Data are shown as mean $\pm$ standard deviation. The normality of the variables was checked with the SHAPIRO-WILK test. To analyze the influence of the intervention program on the study variables, a two-factor ANOVA was performed (Group * pre-post). The significance level was established at $p < 0.05$ in all cases. All calculations were performed with the SPSS version 22 program.

2.5. Ethical Considerations

All families received information about the objectives and characteristics of the study and gave their verbal and written informed consent on behalf of the participant. This work was designed following the deontological standards recognized by the Declaration of Helsinki, complying with the recommendations of the Good Clinical Practice of the Ecuadorian Ministry of Public Health and the current Ecuadorian regulations that regulate research on human beings and received a positive assessment from the Ethics Committee.

3. RESULTS

Table 1: Characteristics of Caregivers.

	Total Sample (N= 200)	GE (N=100)	GC (N=100)	Intergroup difference P
Age (years)	48,25 ± 1,29	47,98 ± 1,21	48,50 ± 1,38	0,35
Years of Care	20,13 ± 1,53	19,83 ± 1,21	20,42 ± 1,77	0,24
Family Functionality	15 ± 2,32	16,5 ± 1,78	13,5 ± 1,78	0,00*
Emotional Well-Being	Somatic symptoms	5,66 ± 0,63	5,83 ± 0,71	0,20
	Anxiety	6 ± 0,65	6,25 ± 0,62	0,06
	Social dysfunction	6,25 ± 0,53	6,33 ± 0,49	0,45
	Depression	5,66 ± 0,86	5,91 ± 0,66	0,16
Caregiver Load (UA)	63,12 ± 4,09	61,6 ± 3,11	64,58 ± 4,56	0,08
Cervical Functionality	55,61 ± 2,23	58,79 ± 2,22	56,83 ± 3,08	0,31

Note: Results expressed as mean ± standard deviation.

GE= Experimental Group; CG = Control Group; P = statistical significance

*indicates a statistically significant intergroup difference with a significance level of P<0.05

The sample was divided into two groups, an intervention group consisting of 100 women aged 47.98 ± 1.21 years, caregivers of patients with CP with a mean age of 19.83 ± 1.21 years of care; and a control group, consisting of 100 women with characteristics similar to the intervention group in age and years of care ($P=0.35$; $P=0.24$). The 2 groups presented mild family dysfunction (13-16 points) with a mean of 15 ± 2.32 points, being lower in the control group compared to the intervention group (13.5 ± 1.78 points vs. 16.5 ± 1.78 points respectively; $P<0.05$).

In the assessment of emotional well-being in the

first data collection, the two groups presented somatic symptoms, anxiety, social dysfunction and depression without intergroup statistical differences ($P>0.05$). Regarding the intensity of the caregiver's burden, the two groups presented an intense burden with a mean of 63.12 ± 4.09 points, presenting intergroup homogeneity ($P=0.08$).

In the assessment of cervical functionality in the first data collection, the two groups presented intense cervical pain, with no statistical differences between groups ($P>0.05$).

Table 2: Characteristics of People with Cerebral Palsy.

	Total Sample (N= 200)	GE (N=100)	GC (N=100)	Intergroup difference P
Age	20.13 ± 1.53 ^a	19.83 ± 1.2 ^a	20.42 ± 1.77 ^a	0,24
Gender	Female	37,50% ^b	16,70% ^b	20,80% ^b
	Male	62,50% ^b	33,30% ^b	29,20% ^b
Total	100% ^b	50% ^b	50% ^b	
Degree of dependence	37.5 ± 10.52 ^a	39.16 ± 11.04 ^a	35.83 ± 10.18 ^a	0,45

Note: a Results expressed as mean ± standard deviation.

b Results expressed in percentages

GE= Experimental Group; CG = Control Group; P = statistical significance

*indicates a statistically significant intergroup difference with a significance level of P<0.05

Regarding the characteristics of the people with CP cared for by the study participants, the male gender predominates with 62.50% of the group of people with CP in the care of the study sample, on the contrary there are 37.50% of women. The average age was 20.13 ± 1.53 years and the degree of dependence among people was moderate, with an average value of 37.5 ± 10.52 score in the Barthel test, presenting intergroup homogeneity ($P>0.05$).

The post-intervention data in the two groups vary intergrouply, in the intervention group the variables

of caregiver burden and emotional well-being decreased statistically significantly ($P<0.05$), on the contrary, in the control group these variables increased.

The caregivers' burden on the Zarit test was significantly reduced in the group that followed the $P<0.05$ exercise program. On the other hand, in the control group there was no statistically significant change ($P=0.08$).

When analysing emotional well-being, the subscales that showed the greatest reduction after

the intervention with the physical activity programme were anxiety and depression ($P < 0.05$), while the control group statistically significantly increased anxiety ($P = 0.01$) and social dysfunction ($P = 0.03$).

The experimental group reduced neck pain after the exercise intervention ($P = 0.01$), unlike the control group, which did not present any statistically significant change ($P = 0.08$).

Table 3: Caregiver Burden, Emotional Well-Being, and Neck Pain Before and After Surgery.

		GE (N=100)	Intragroup Difference P	GC (N=100)	Intragroup Difference P	Intergroup Difference P
Caregiver Burden	pre	61,6 ± 3,11	0,00*	64,58 ± 4,56	0,08	0,00*
	post	53,6 ± 3,72		65,25 ± 4,15		
Somatic symptoms	pre	5,83 ± 0,71	0,00*	5,5 ± 0,52	0,08	0,015*
	post	4,9 ± 0,79		5,75 ± 0,75		
Anxiety	pre	6,25 ± 0,62	0,00*	5,75 ± 0,62	0,01*	0,00*
	post	4 ± 0,6		6,16 ± 0,83		
Social dysfunction	pre	6,33 ± 0,49	0,00*	6,16 ± 0,57	0,03*	0,00*
	post	5,08 ± 0,66		6,5 ± 0,67		
Depression	pre	5,91 ± 0,66	0,00*	5,41 ± 0,99	0,08	0,012*
	post	4,08 ± 0,79		5,16 ± 1,11		
Cervicalgia	pre	58,79 ± 2,22	0,00*	56,83 ± 3,08	0,08	0,012*
	post	44,08 ± 1,79		57,16 ± 2,41		

Note: GE= Experimental group; CG = Control Group; P = statistical significance. *Indicates a statistically significant intragroup difference with a significance level $P < 0.05$

4. DISCUSSION

The physical exercise program applied in Caregiver Overload Syndrome produced positive effects in decreasing caregiver burden, somatic symptoms, anxiety, social dysfunction, and depression in the intervention group, which unlike the control group, at the end of 16 weeks increased anxiety and social dysfunction.

The literature mentions that practicing physical exercise in a group expands and improves social interaction, due to the development of social, communicative and co-existence skills and abilities through body expression, cooperation and inclusion. In addition, physical exercise improves people's self-esteem, confidence, commitment, effort and vitality by feeling better about themselves, which favors interpersonal relationships. (11-15)(16-18)

The results of this research agree with studies carried out in caregivers of people with chronic diseases, that the practice of physical exercise causes a positive anxiolytic effect and the reduction of symptoms of depression and anxiety. Physical exercise improves the regulation of the cardiovascular and respiratory system by affecting the nervous system, which increases the levels of neurotransmitters such as: endorphins, noradrenaline, serotonin, reducing anxiety, pain and improving mood and the feeling of well-being. (19,20)(21,22).

The results of the present study collaborate as a

non-pharmaceutical treatment for anxiety and neck pain, with physical exercise for 16 weeks being a good option to alleviate the symptoms of Caregiver Overload Syndrome.

5. CONCLUSION

In caregivers of people with chronic diseases, such as CP, they usually present exhaustion, physical and psychological exhaustion due to intense dedication to care. This situation causes a series of physical, psychological and emotional symptoms.

All the aforementioned symptomatology treated with the intervention of a physical exercise program for 16 weeks in an experimental group of 100 informal caregivers of CP improved caregiver burden, emotional health and cervical functionality. In contrast, the control group after 16 weeks worsened emotional health in the aspects of anxiety and social dysfunction.

Physical exercise improved physical and emotional health, due to the development of social and communicative skills and abilities through body expression, also the positive effects of physical exercise were to regulate the nervous system, increasing the release of neurotransmitters improving mood and the feeling of well-being.

This study is of great contribution to the treatment of neck pain and anxiety with physical exercise for 16 weeks, good results are observed in the research sample.

REFERENCES

- Duvignau E. Quality of life in patients. *Rev Sanid Milit Mex.* [Inter-net]. 2015; 69(4):535–42. Available at: <https://www.medigraphic.com/pdfs/sanmil/sm-2015/sm156e.pdf>.
- Garip Y. Fatigue in the mothers of children with cerebral palsy. *Disability and Rehabilitation.* 2017; 39(8):757–62. doi: 10.3109/09638288.2016.1161837.
- Chamorro A. Characterization of the Primary Caregiver of Patients with Neurological Disabilities. *Journal of the Faculty of Health Sciences of the University of Cauca* [Internet]. 2009; 17(3):31–8. Available at: <https://revistas.unicauca.edu.co/index.php/rfcs/article/view/214>.
- Lamotte G. Exercise Training for Persons with Alzheimer's Disease and Caregivers: A Review of Dyadic Exercise Interventions. *Journal of motor behavior.* 2017; 49(4):365–77. doi: 10.1080/00222895.2016.1241739.
- Ghaderi F, Mohammadi K, Amir Sasan R, Niko Kheslat S, Oskouei AE. Effects of Stabilization Exercises Focusing on Pelvic Floor Muscles on Low Back Pain and Urinary Incontinence in Women. *Urology* [Internet]. 2016; 93:50–4. Disponible en: <https://pubmed.ncbi.nlm.nih.gov/27059833/>.
- Cordeiro LMS, Rabelo PCR, Moraes MM, Teixeira-Coelho F, Coimbra CC, Wanner SP, et al. Physical exercise-induced fatigue: the role of serotonergic and dopaminergic systems. *Braz J Med Biol Res.* 2017; 50(12):e6432. doi: 10.1590/1414-431X20176432. PubMed PMID: 29069229.
- Connell CJW, Thompson B, Turuwhenua J, Szlachetka A, Gant N. Effects of Dopamine and Norepinephrine on Exercise-induced Oculomotor Fatigue. *Medicine and science in sports and exercise* [Inter-net]. 2017; 49(9):1778–88. Disponible en: <https://pubmed.ncbi.nlm.nih.gov/28452866/>.
- Carek P. Exercise for the treatment of depression and anxiety. *Int J Psychiatry Med.* 2011; 41(1):15–28. doi: 10.2190/PM.41.1.
- Cuthbert CA, King-Shier K, Tapp DM, Ruether D, Jackson C, Culos-Reed SN. Renewing caregiver health and wellbeing through exercise (RECHARGE): A randomized controlled trial. *Contemp Clin Trials.* 2016; 50:273–83. doi: 10.1016/j.cct.2016.08.007. PubMed PMID: 27530087.
- Botha E, Gwin T, Purpora C. The effectiveness of mindfulness based programs in reducing stress experienced by nurses in adult hospital settings: a systematic review of quantitative evidence protocol. *JBIS Database System Rev Implement Rep.* 2015; 13(10):21–9. doi: 10.11124/jbisr-2015-2380. PubMed PMID: 26571279.
- Thompson D, Karpe F, Lafontan M, Frayn K. Physical activity and exercise in the regulation of human adipose tissue physiology. *Physiol Rev.* 2012; 92(1):157–91. doi: 10.1152/physrev.00012.2011. PubMed PMID: 22298655.
- Yu F, Thomas W, Nelson NW, Bronas UG, Dysken M, Wyman JF. Impact of 6-month aerobic exercise on Alzheimer's symptoms. *J Appl Gerontol.* 2015; 34(4):484–500. doi: 10.1177/0733464813512895. PubMed PMID: 24652914.
- Wegner M. Effects of exercise on anxiety and depression disorders: review of meta-analyses and neurobiological mechanisms. *CNS & neurological disorders drug targets.* 2014; 18(7). doi: 10.2174/1871527313666140612102841.
- Del Cura Bilbao A, Sandín Vázquez M. Assets for health and quality of life in people diagnosed with serious mental illness. *Gac Sanit.* 2020. doi: 10.1016/j.gaceta.2020.03.004. PubMed PMID: 32467001.
- Cuthbert CA, King-Shier KM, Ruether JD, Tapp DM, Wytsma-Fisher K, Fung TS, et al. The Effects of Exercise on Physical and Psychological Outcomes in Cancer Caregivers: Results From the RECHARGE Randomized Controlled Trial. *Ann Behav Med.* 2018; 52(8):645–61. doi: 10.1093/abm/kax040. PubMed PMID: 30010704.
- Orgeta V, Miranda-Castillo C. Does physical activity reduce burden in carers of people with dementia? A literature review. *International journal of geriatric psychiatry* [Internet]. 2014; 29(8):771–83. Disponible en: <https://pubmed.ncbi.nlm.nih.gov/25191688/>.
- Stanton R. Exercise and the treatment of depression: a review of the exercise program variables. *Journal of science and medicine in sport* [Internet]. 2014; 28(3):177–82. Disponible en: <https://doi.org/10.1016/j.jsams.2013.03.010>.
- Singh N. The efficacy of exercise as a long-term antidepressant in elderly subjects: a randomized, controlled trial. *The journals of gerontology.* 2001; 56(8):497–504. doi: 10.1093/gerona/56.8.m497.
- Morland C, Andersson KA, Haugen ØP, Hadzic A, Kleppa L, Gille A, et al. Exercise induces cerebral VEGF and

- angiogenesis via the lactate receptor HCAR1. *Nat Commun.* 2017;8:15557. doi: 10.1038/ncomms15557. PubMed PMID: 28534495.
- Ochentel O. Efficacy of Exercise Therapy in Persons with Burnout. A Systematic Review and Meta-Analysis. *Journal of sports science & medicine* [Internet]. 2018; 16(1):475–84. Disponible en: https://pubmed.ncbi.nlm.nih.gov/30116121-efficacy-of-exercise-therapy-in-persons-with-burnout-a-systematic-review-and-meta-analysis/?from_term=exercise+and+burnout+&from_pos=1.
- Lowery D. The effect of exercise on behavioural and psychological symptoms of dementia: the EVIDENCY. 2014.
- Madrugá M, Prieto J, Rohlfes P, Gusi N. Cost-Effectiveness and Effects of a Home-Based Exercise Intervention for Female Caregivers of Relatives with Dementia: Study Protocol for a Randomized Controlled Trial. *Healthcare (Basel)*. 2020; 8(1). doi: 10.3390/healthcare8010054. PubMed PMID: 32155761.