

DOI: 10.5281/zenodo.12426507

THE IMPACT OF HEALTHCARE WORKER BURNOUT AND FATIGUE ON PATIENT SAFETY: A MULTIDISCIPLINARY ANALYSIS

Amal Ali Alghorabi^{1*}, Wejdan Mohammed Al-Mutairi², Huda Ali Alqahtani³, Sarah Abdullah al Manea⁴, Munirah Saeed Alqahtani⁵, Hameedah Mansi Al_Alawi⁶, Essa Hamza ALamri⁷, Ghaya Mansour Fatani⁸, Jaber Omar Jaber Asiri⁹, Najla yousef alsanea¹⁰

^{1 3 6 8} Pharmacist, Health Services of the Ministry of Defense, King Abdulaziz Airbase Hospital, Dhahran, Saudi Arabia

^{2 7 9} Pharmacy Technician, Health Services of the Ministry of Defense, King Abdulaziz Airbase Hospital, Dhahran, Saudi Arabia

⁴ Senior Pharmacist, Health Services of the Ministry of Defense, King Abdulaziz Airbase Hospital, Dhahran, Saudi Arabia

⁵ Dietitian, Health Services of the Ministry of Defense, King Abdulaziz Airbase Hospital, Dhahran, Saudi Arabia

¹⁰ Nurse, Health Services of the Ministry of Defense, King Abdulaziz Airbase Hospital, Dhahran, Saudi Arabia

Received: 11/11/2025

Accepted: 07/03/2026

Corresponding author: Amal Ali Alghorabi

[\(Email\)](#)

ABSTRACT

Patient safety is fundamentally linked to the wellbeing of the healthcare workforce, yet systemic issues like burnout and fatigue continue to jeopardize clinical outcomes and health system stability. This article examines the intersection of provider exhaustion and patient harm, reframing burnout as a signal of misaligned system design rather than an individual failing. This synthesis draws upon a cross-sectional study of internal medicine residents⁶, a quality improvement evaluation of an AI-driven scheduling system in anesthesiology, a global situational analysis from the World Health Organization, a meta-analysis of organizational interventions, a four-year longitudinal study of overtime in Taiwan¹⁰, and a mixed-method clinical audit of sleep deprivation in the UK. Dramatic changes to scheduling models have been shown to cause high levels of fatigue and a much higher risk to both personal and occupational health. Studies indicate that traditional "scheduling models" (for example, the 1-in-4 resident schedule) have the highest levels of fatigue and very high levels of occupational burn out (HR = 2.13), with Nurses and Physicians being at a much higher risk than any other staff in the organization. Long-term data suggest that physicians and nurses are also at a much higher risk for developing occupational burn out than the general population. Conversely, when Structural interventions are used to increase access to AI-driven scheduling methods, Intraoperative Care Processes are reduced by 9.1%. The estimated cost of associated harm incurred from this 9.1% will be \$335,550. There also were multiple studies that have confirmed that for every additional patient added to a Nurse, there is a 7% increase in the risk of death from hospital-associated complications. In order to address the patient safety crisis, it will be necessary to move away from individual coping strategies to a system of accountability. It will be necessary to implement alternative scheduling models, mandated staffing ratios, and to implement a "just culture." Building high-reliability health systems will require legal protections for incident reporting.

Keywords: Healthcare Worker, Burnout, Fatigue, Patient Safety.

1. INTRODUCTION

1.1. The Intersection of Fatigue and Harm

This pursuit of UHC is a laudable goal, but it is also fundamentally premised on ensuring health service safety (World Health Organization [WHO, 2024]). In the absence of this, UHC translates into a lack of trust and a reluctance to seek healthcare (WHO, 2024). Patient safety concerns are presently costing health systems billions of dollars in costs, with a staggering 12.6% of total health expenditure in high-income countries (HICs) being spent on managing harm (WHO, 2024). At the heart of this health safety crisis is a two-pronged epidemic of healthcare worker burnout and fatigue, now recognized as work-related hazards as opposed to personal failings (Yuan et al., 2023; Zavaleta - Monestel et al., 2026).

1.1.1. Global Burden and Economic Cost of Unsafe Care

Preventable harm in healthcare is not only a health concern; it is also a massive economic burden (WHO, 2024). For OECD countries, managing harm in healthcare translates into a staggering US\$ 878 billion in annual costs (WHO, 2024) In middle-income countries such as Thailand, preventable harm in healthcare occurs in 7% of healthcare admissions, translating into a staggering 5.5% of the national health budget (WHO, 2024).

1.2. Burnout: A Systemic Design Flaw

However, burnout is now being seen less as a failure of personal resilience but more as an indicator of poorly designed systems (Zavaleta-Monestel et al., 2026). Burnout is marked by emotional exhaustion, depersonalization, and reduced feelings of personal accomplishment (Bes et al., 2023; Yuan et al., 2023; Zavaleta-Monestel et al., 2026). Studies have revealed that workload, technology, and management are the major causes of this condition (Zavaleta-Monestel et al., 2026). In the pharmacy industry, for example, the system is largely designed for throughput rather than sustainability, emphasizing speed over other forms of cognitive work such as patient counseling and clinical decision-making (Zavaleta-Monestel et al., 2026). This is the major reason for the systematic devaluation of cognitive work, which leads to

burnout (Zavaleta-Monestel et al., 2026). Longitudinal research has revealed that working long hours is a major factor that leads to burnout. A study revealed that workers who work long hours are at a 113% greater risk of suffering from high burnout compared to those working fewer hours (Chen et al., 2025). Females, physicians, and nurses are more prone to burnout due to work-home conflicts, high-stress patient interactions, and other reasons (Chen et al., 2025).

1.3. The Critical Role of Sleep and Fatigue Management

Fatigue is one of the major causes of medical errors and injuries (WHO, 2024; Chukwunonso-Ogbu et al., 2025). Prolonged wakeful periods for 24 hours impair cognitive function to the extent that the person is considered to be driving under the influence of alcohol, exceeding the permissible limit (Yuan et al., 2023; Chukwunonso-Ogbu et al., 2025). In critical health-care environments, it was found that more than half of the staff had less than six hours of sleep before the start of work, and 88% of the staff worked while fatigued (Chukwunonso-Ogbu et al., 2025).

The consequences of this sleep deprivation are severe. They are as follows:

1. Cognitive Lapses: Difficulty in concentrating, forgetting medications, and delayed response times are reported by the staff (Chukwunonso-Ogbu et al., 2025).
2. Physical Hazards: Fatigue may lead to occupational injuries, household injuries, and management of personal health (Yuan et al., 2023).
3. Normalization of Exhaustion: Exhaustion is a norm among the health-care staff, which may lead to the underreporting of errors and reluctance to speak up for safety concerns (Chukwunonso-Ogbu et al., 2025).

WHO 7x5 Matrix for Patient Safety offers a guide for action, with a focus on education for health workers, patient participation, and risk management. Education for undergraduate and postgraduate students about patient safety is a necessity but a challenge for only 20% of the countries to implement effectively.

Table 1: Comparative Analysis of Staffing and Scheduling Impacts

Feature	Traditional 1-in-4 Model	Night Float/Shift-Based	AI-Driven Scheduling	Mandated Staffing Ratios
Shift Duration	Exceeds 24 hours	8-12 hours	Optimized 8-13 hours	Varied based on unit
Burnout Rate	Highest (71.4%)	Moderate (57.1%)	Significantly Reduced	24% Lower Odds
Safety Impact	High risk of medical error	Lower fatigue hazards	9.1% less care transitions	7% lower mortality per patient

Primary Driver	Tradition/Chronicity	Shift-based coverage	Combinatorial optimization	Legislated safety floor
Wellbeing Focus	Personal challenge	Protected personal health	Flexibility/Preference	Workforce retention

1.4. From Individual Coping to System Accountability

In the modern healthcare system, the need to change the focus from individual endurance to system accountability is a necessity. Interventions at the organizational level to manage workload, scheduling, and staffing ratios not only work for preventing burnout but also prove to be cost-effective by offering a high ROI.

In order to ensure patient safety, the following need to be taken care of:

1. Explore alternative work patterns like night float shifts, as well as the use of artificial intelligence technology, to mitigate fatigue-related hazards.
2. Ensure the implementation of the mandatory staffing ratios to protect both nurses and patients, thereby ensuring the "surveillance system" of nursing is maintained.
3. Cultivate a Just Culture where openness and reporting of incidents are promoted, along with legal protections for health workers.
4. Invest in rest facilities, such as sleep pods, to counter the cognitive effects of the inevitable shift work required.

Through this realization that "if it's not safe, it's not care," political leaders and medical professionals can unite to transform the delivery of healthcare into a sustainable high-reliability system.

2. METHODS

The research synthesized in this article takes a multi-dimensional approach to the study of workforce wellbeing and safety:

1. Cross-Sectional Surveys: This approach surveyed internal medicine residents (PGY-1 to PGY-3) to identify the effects of the impact of five different scheduling models on fatigue and burnout (Yuan et al., 2023). Burnout was surveyed through the Maslach Burnout Inventory (MBI), while fatigue was surveyed through the Swedish Occupational Fatigue Inventory (SOFI) (Yuan et al., 2023).
2. Longitudinal Survival Analysis: This approach surveyed a four-year study in Taiwan (2021-2024) to identify the effects of long hours of overtime work on the progression to high levels of personal burnout through the Kaplan Meier

method and Cox regression analysis (Chen et al., 2025).

3. Quality Improvement Evaluation: This approach surveyed the effects of the impact of the AI-based scheduling system Lightning Bolt through the idealized design framework on the burnout of anesthesiologists (Sumrall et al., 2025). Metrics surveyed include intraoperative transitions of care, physician satisfaction, and vacation approval rates (Sumrall et al., 2025).
4. Meta-Analysis: This approach surveyed the effectiveness of organizational interventions versus combined interventions to reduce exhaustion scores through the analysis of 13 studies (Bes et al., 2023).
5. Clinical Audit: A mixed-method audit in high-acuity UK hospitals, using anonymous surveys and face-to-face interviews, found sleep deprivation and its impact on performance (Chukwunonso-Ogbu et al., 2025).
6. Global Policy Review: WHO's Global Patient Safety Report 2024 employed a structured survey of 108 Member States to assess the achievement of a 7x5 strategic matrix of safety goals (WHO, 2024).

3. RESULTS

The study's results prove that systemic factors are the main contributors to provider exhaustion and patient harm. In residency training, the classic 1-in-4 call model, with 24-hour shifts every four days, showed the highest levels of burnout (71.4%) and a substantial increase in risks to personal health and occupational harm (Yuan et al., 2023). Longitudinal studies found that workers experiencing overtime are 113% more likely to experience high levels of burnout than those not experiencing overtime (Chen et al., 2025).

Technological and policy interventions showed clear benefits. The implementation of AI scheduling led to a 9.1% reduction in transitions of care and a massive 82% decrease in vacation denial rates (Sumrall et al., 2025). In nursing, mandated staffing ratios in Queensland resulted in 145 fewer deaths and a return on investment where savings from reduced readmissions were double the cost of additional staff (Aiken, 2025).

Table 2: Burnout Prevalence and Risk Factors by Demographic and Role

Variable	Findings/ Risk Ratio	Context/ Source
Overall Burnout Prevalence	58%	Residents (Yuan et al., 2023)
Overtime Work	HR 2.13 (113% increase)	Hospital Workers (Chen et al., 2025)
Physician Role	HR 1.79 (79% increase)	Higher risk vs. other staff (Chen et al., 2025)

Nurse Role	HR 1.57 (57% increase)	Higher risk vs. other staff (Chen et al., 2025)
Female Gender	HR 1.49 (49% increase)	Gender disparity (Chen et al., 2025)
Reduced Sleep	HR 1.29 (29% increase)	Independent risk factor (Chen et al., 2025)

Table 3: Impact of Scheduling Models and Fatigue on Safety and Wellbeing

Metric	Traditional 1-in-4 Model	AI-Driven/ Alternative	Source
Burnout Rate	71.4% (Highest)	53.3% (Late-stay)	(Yuan et al., 2023)
Occupational Harm Risk	OR 5.69	Not significant	(Yuan et al., 2023)
Care Transitions	10.3% (Manual)	9.1% (AI-Driven)	(Sumrall et al., 2025)
Vacation Denial Rate	N/A	82% Decrease	(Sumrall et al., 2025)
Worked While Fatigued	88% of staff	N/A	(Chukwunonso-Ogbu, 2025)
Sleep Before Shift	< 6 hours (50% of staff)	N/A	(Chukwunonso-Ogbu, 2025)

Table 4: Clinical and Economic Impact of Nurse Staffing Ratios

Jurisdiction	Staffing Policy Impact	Economic / Safety Outcome	Source
Global Standard	Each +1 patient per nurse	7% increase in mortality risk	(Aiken, 2025)
Queensland, AU	Implemented Ratios	145 fewer deaths; 255 fewer readmissions	(Aiken, 2025)
Queensland, AU	AUD \$70M Savings	2x the cost of additional staff	(Aiken, 2025)
California, US	Implemented Ratios	Reduced odds of poor quality care (42%)	(Aiken, 2025)
Illinois, US (Est)	Proposed 1:4 ratio	1,595 deaths avoided; \$117M saved	(Aiken, 2025)
New York (Est)	Proposed 1:4 ratio	More sepsis deaths avoided than care bundles	(Aiken, 2025)





Figure 1: Patient safety in national health budgets The scarcity of patient safety human resource plans and ongoing efforts to address staffing gaps emphasizes the urgent need for comprehensive strategies to ensure there are sufficient health workers.



Figure




Figure 2: Number of countries that responded to the Member State survey, by World Bank income country classification 2022-2023




Figure 3: Proportion of countries that have defined minimum safety standards Global integration of safety standards into health care licensing emphasizes a commitment to universal safety norms

4. DISCUSSION

4.1. Reframing Burnout as a Systemic Design Flaw

The synthesis of current research confirms that exhaustion is the core dimension of burnout and serves as a primary signal that a healthcare system's design is misaligned (Bes et al., 2023; Zavaleta-Monestel et al., 2026). Burnout is increasingly recognized not as a lack of individual resilience, but as a predictable consequence of structural imbalances where workload demands consistently exceed available resources (Zavaleta-Monestel et al., 2026). In sectors such as pharmacy, systems are often structured for throughput (volume) rather than sustainability, prioritizing speed over essential cognitive labor like clinical reasoning and patient counseling (Zavaleta-Monestel et al., 2026). This systematic undervaluation of professional expertise accelerates disengagement and emotional fatigue (Zavaleta-Monestel et al., 2026).

The Dangerous Normalization of Fatigue Clinical evidence reveals a pervasive "normalization of exhaustion" within hospital cultures, where working while sleep-deprived is often viewed as a personal challenge rather than an occupational hazard (Yuan et al., 2023; Chukwunonso-Ogbu et al., 2025). A clinical audit found that 88% of staff had worked while fatigued, and nearly 40% confirmed witnessing or making errors where fatigue was a contributing factor (Chukwunonso-Ogbu et al., 2025). This is particularly alarming given that 24 hours of sustained wakefulness produces cognitive impairments equivalent to being legally intoxicated (Yuan et al., 2023). Despite these risks, many resident physicians still do not conceptualize fatigue as a threat to patient safety, which underscores the urgent need for context-specific fatigue risk management education (Yuan et al., 2023). This systematic undervaluing of professional expertise contributes to disengagement and emotional fatigue (Zavaleta-Monestel et al., 2026).

The Dangerous Normalization of Fatigue Clinical evidence confirms the normalization of exhaustion as a dominant phenomenon in the cultures of hospitals, where sleep deprivation while working is perceived as a personal challenge rather than an occupational hazard (Yuan et al., 2023; Chukwunonso-Ogbu et al., 2025). A clinical audit reported that 88% of the staff reported working while fatigued, with nearly 40% admitting to having witnessed or committed errors due to fatigue (Chukwunonso-Ogbu et al., 2025). This phenomenon is alarming since 24 hours of wakefulness results in cognitive impairments similar

to those of a person who is legally intoxicated (Yuan et al., 2023). However, most resident physicians do not recognize fatigue as a patient safety threat, a fact that highlights the need for fatigue risk management education to be context-specific (Yuan et al., 2023).

4.2. The Direct Link Between Staffing, Overtime, and Mortality

Longitudinal data and policy reviews establish a clear, causal link between staffing levels and patient survival. In Taiwan, a four-year study found that healthcare workers experiencing prolonged overtime faced a 113% higher risk of progressing to high personal burnout (Chen et al., 2025). Similarly, landmark nursing research has documented that each additional patient added to a nurse's workload increases the risk of hospital mortality by 7% (Aiken, 2025). These findings suggest that nurses serve as a critical "surveillance system" for detecting complications; when staffing ratios are too high, this system fails, leading to life-threatening "failure to rescue" events (Aiken, 2025).

Organizational Accountability vs. Individual Coping The sources argue for a paradigm shift from individual-focused wellness initiatives to organizational accountability (Zavaleta-Monestel et al., 2026). While many health systems offer resilience training or wellness apps, meta-analyses indicate that workload-focused interventions—such as increasing human resources or optimizing workflows—are significantly more effective at reducing exhaustion scores (Bes et al., 2023). Structural reforms, such as the implementation of AI-driven scheduling, have shown the ability to reduce hazardous intraoperative handoffs by 9.1% while simultaneously decreasing vacation denial rates by 82% (Sumrall et al., 2025). This demonstrates that technological innovation can align physician satisfaction with patient safety goals (Sumrall et al., 2025). This shows that technological innovation can actually align physician satisfaction with patient safety goals (Sumrall et al., 2025).

Building High Reliability Systems through "Just Culture" To build high reliability health systems, a global commitment to "just culture" is needed, where learning from mistakes is balanced with professional accountability (WHO, 2024). Currently, a major gap exists in this area because, although 55% of countries worldwide recognize patient safety as a national priority, only 19% have laws in place to protect health workers who report patient safety incidents (WHO, 2024). As a consequence, a culture of fear exists, preventing openness in learning from mistakes (WHO, 2024). Furthermore, the WHO points out that only 11% of countries have sufficient

financial and human resources to support their patient safety strategies, thus a political commitment is needed to ensure "if it's not safe, it's not care" (WHO, 2024).

5. CONCLUSION

However, patient safety cannot be disassociated from the safety of the workforce, as emphasized by the WHO's Director-General, "if it's not safe, it's not care". The 1-in-4 resident call model, as well as the chronic understaffing of nurses, is recognized as an occupational hazard that directly leads to increased mortality and error rates. To address this crisis, health organizations should adopt alternative work schedules such as the night float schedule or the AI

schedule, which is optimized for the workforce's sleep patterns.

Organizations should shift the focus from the workforce's resilience to organizational accountability, such as the implementation of the legislated floor for staffing. Moreover, the development of a just culture with legal protections for reporting is critical for identifying the root causes of the harm. By investing in the workforce through the appropriate allocation of resources, health organizations can achieve the dual purpose of sustainable professional practice and the complete elimination of avoidable patient harm.

REFERENCES

1. Yuan, J. H., Huang, Y., Rosgen, B. K., Donnelly, S., Lan, X., & Katz, S. J. (2023). Burnout and fatigue amongst internal medicine residents: A cross-sectional study on the impact of alternative scheduling models on resident wellness. *PLOS ONE*, 18(9), e02914571.
2. Sumrall, W. D., Oury, J. V., & Gilly, G. M. (2025). Enhancing Physician Satisfaction and Patient Safety Through an Artificial Intelligence-Driven Scheduling System in Anesthesiology. *Ochsner Journal*, 25(1), 44-492.
3. World Health Organization. (2024). *Global Patient Safety Report 2024*. Geneva: World Health Organization. ISBN 978-92-4-009545-834.
4. Bes, I., Shoman, Y., Al-Gobari, M., Rousson, V., & Guseva Canu, I. (2023). Organizational interventions and occupational burnout: a meta-analysis with focus on exhaustion. *International Archives of Occupational and Environmental Health*, 96(9), 1211-12235.
5. Zavaleta-Monestel, E., Söderlund, L. Å., Guiu-Segura, J. M., & Arguedas-Chacón, S. (2026). Pharmacist burnout: from coping to system accountability in the medication-use process. *Frontiers in Public Health*, 13, 174933267.
6. Chen, Y.-H., Jong, G.-P., Yang, C.-W., & Lee, C.-H. (2025). Prolonged Overtime Predicts Worsening Burnout among Healthcare Workers: A 4-Year Longitudinal Study in Taiwan. *Preprints.org*. doi: 10.20944/preprints202506.2060.v189.
7. Chukwunonso-Ogbu, A., Fazli, S. A., Kalungi, G., & Malomo, O. (2025). Sleep Deprivation and Fatigue in Healthcare Staff: A Clinical Audit on the Risk to Patient Safety. *Cureus*, 17(11), e965431011.
8. Aiken, L. H. (2025). *Staffing Ratios and their Impact on the Health and Safety of Nurses: A Policy Brief*. Prepared for the Ontario Nurses Association12.
9. McHugh, M. D., Aiken, L. H., Sloane, D. M., Windsor, C., Douglas, C., & Yates, P. (2021). Nurse staffing and patient mortality, readmissions, and length of stay: A prospective study of the effects of nurse-to-patient ratio legislation in a panel of hospitals. *The Lancet*, 397, 1905-191313.
10. Slawomirski, L., & Klazinga, N. (2022). The economics of patient safety: from analysis to action. *OECD Health Working Papers* No. 1451415.
11. Needleman, J., Buerhaus, P. I., Mattke, S., Stewart, M., & Zelevinsky, K. (2002). Nurse staffing levels and quality of care in hospitals. *New England Journal of Medicine*, 346(22), 1715-172216.
12. Dawson, D., & Reid, K. (1997). Fatigue, alcohol and performance impairment. *Nature*, 388(6639), 23517.
13. Barger, L. K., Weaver, M. D., Sullivan, J. P., Qadri, S., Landrigan, C. P., & Czeisler, C. A. (2023). Impact of work schedules of senior resident physicians on patient and resident physician safety: nationwide, prospective cohort study. *BMJ Medicine*, 2(1), e00032018.
14. Jha, A. K., Larizgoitia, I., Audera-Lopez, C., Prasopa-Plaizier, N., Waters, H., & Bates, D. W. (2013). The global burden of unsafe medical care: analytic modelling of observational studies. *BMJ Quality & Safety*, 22, 809-81519.
15. Rotenstein, L. S., Torre, M., Ramos, M. A., Rosales, R. C., Guille, C., Sen, S., et al. (2018). Prevalence of Burnout Among Physicians. *JAMA*, 320(11), 113120.

16. Maslach, C., & Jackson, S. E. (1981). The measurement of experienced burnout. *Journal of Organizational Behavior*, 2(2), 99-11320.
17. Kecklund, G., & Axelsson, J. (2016). Health Consequences of Shift Work and Insufficient Sleep. *BMJ*, 355, i521021.
18. Lasater, K. B., Aiken, L. H., Sloane, D. M., French, R., Anusiewicz, C. V., Martin, B., et al. (2021). Is hospital nurse staffing legislation in the public's interest? An observational study in New York State. *Medical Care*, 59(5), 444-45022.
19. Shanafelt, T. D., Balch, C. M., Bechamps, G., Russell, T., Dyrbye, L., Satele, D., et al. (2010). Burnout and Medical Errors Among American Surgeons. *Annals of Surgery*, 251(6), 995-10002324.
20. Thomas, A. D., Pandit, C., & Krevat, S. A. (2020). Race differences in reported harmful patient safety events in healthcare system high reliability organizations. *Journal of Patient Safety*, 16, e235-e23925.